

Weight loss and bone mineral content

Citation for published version (APA):

Westerterp, K. R. (2002). Weight loss and bone mineral content. *Obesity Research*, 10(6), 559.
<https://doi.org/10.1038/oby.2002.76>

Document status and date:

Published: 01/01/2002

DOI:

[10.1038/oby.2002.76](https://doi.org/10.1038/oby.2002.76)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Weight Loss and Bone Mineral Content

Klaas R. Westerterp

Weight loss is accompanied by a decrease in bone mineral content. The main explanation is probably a reduction of the mechanical strain on the skeleton. Thus, the addition of exercise to weight-loss programs may reduce the risk for bone loss (1). However, exercise training during energy restriction seems to be compensated by a reduction of activity outside of training sessions (2,3). A diet intervention to reduce the weight-loss induced decrease in bone mineral content could be more appropriate. The most straightforward intervention is calcium supplementation. Indeed calcium supplementation, even if the diet contains the officially recommended amounts of calcium, partly inhibits weight loss accompanied bone loss (4). Interestingly, Skov et al. (5) show a similar protective effect of protein intake on weight-loss induced bone loss.

Protein intake has been shown to be important in maintaining bone or minimizing bone loss in elderly persons (6). After controlling for weight loss, women and men with a relatively higher protein intake had decreased bone loss. Initially, it was thought that a high protein intake caused bone loss instead of protection against bone loss. A higher dietary protein intake leads to higher urinary calcium excretion. The increase in urinary calcium was a function of the amount and type of protein intake because dietary acid load is slightly influenced by the ratio of meat to vegetable protein intake (7). Later studies showed that total calcium balance is positively affected by nitrogen intake despite the higher urinary calcium excretion because of a lower calcium excretion in feces (8). The study of Skov et al. (5) reflects the positive effect of nitrogen on calcium balance in bone mass.

The conclusion of the study of Skov et al. (5) is that body-fat loss was the major determinant of loss of bone mineral content. Unfortunately, no mechanism for the relation was mentioned or discussed. The mediator between

body fat and bone may be leptin. Serum leptin concentrations have been shown to be associated with bone mass (9). Leptin stimulates mineralization of the bone matrix. A reduction of body fat could affect bone mass through the effect of fat mass on serum leptin concentrations.

In conclusion, the effect of weight loss on bone mass is 2-fold: a reduction of the mechanical strain on the skeleton and a reduction through fat loss with leptin as the potential mediator. Nitrogen intake can partly reduce the effect through a positive effect on net calcium balance.

References

1. **Ryan AS, Nicklas BJ, Dennis KE.** Aerobic exercise maintains regional bone mineral density during weight loss in postmenopausal women. *J Appl Physiol.* 1998;84:1305–10.
2. **Racette SB, Schoeller DA, Kushner RF, Neil KM, Herling-Iaffaldano K.** Effects of aerobic exercise and dietary carbohydrate on energy expenditure and body composition during weight reduction in obese women. *Am J Clin Nutr.* 1995;61:486–94.
3. **Kempen KPG, Saris WHM, Westerterp KR.** Energy balance during 8 weeks energy-restrictive diet with and without exercise in obese females. *Am J Clin Nutr.* 1995;62:722–9.
4. **Jensen LB, Kolerup G, Quaade F, Sorensen OH.** Bone minerals changes in obese women during a moderate weight loss with and without calcium supplementation. *J Bone Miner Res.* 2001;16:141–7.
5. **Skov AR, Haulrik N, Toubro S, Molgaard C, Astrup A.** The effect of protein intake on bone mineralization during weight loss: a randomised controlled 6-months trial in overweight subjects. *Obes Res.* 2002;10:432–8.
6. **Hannan MT, Tucker KL, Dawson-Hughes B, Cuppes LA, Felson DT, Kiel DP.** Effect of dietary protein on bone loss in elderly men and women: the Framingham Osteoporosis Study. *J Bone Miner Res.* 2000;15:2504–12.
7. **Heaney RP.** Nutritional factors in osteoporosis. *Annu Rev Nutr.* 1993;13:287–316.
8. **Pannemans DLE., Schaafsma GJ, Westerterp KR.** Calcium excretion, apparent calcium absorption and calcium balance in young and elderly subjects: influence of protein intake. *Br J Nutr.* 1997;77 721–9.
9. **Pasco JA, Henry MJ, Kotowicz MA, Collier GR, Ball MJ, Ugoni AM, Nicholson GC.** Serum leptin levels are associated with bone mass in non-obese women. *J Clin Endocrinol Metab.* 2001;86:1884–7.

Submitted for publication. Accepted in final form. Department of Human Biology, Maastricht University, Maastricht, The Netherlands.

Address correspondence to, Klaas R. Westerterp, Department of Human Biology, Maastricht University, PO Box 616, 6200 MD Maastricht, The Netherlands.

E-mail: K.Westerterp@HB.unimaas.nl

Copyright © 2002 NAASO